Un nuovo anticorpo anti-CD19

Pier Luigi Zinzani Institute of Hematology «L. e A. Seràgnoli» University of Bologna

Disclosures (1): PIER LUIGI ZINZANI

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other
Secura Bio						x	
Celltrion					x	x	
Gilead					x	x	
Janssen-Cilag					x	x	
BMS					x	x	
Servier					x	x	
Sandoz						x	
MSD			x		x	x	
TG Therap.					х	X	

Disclosures (2): PIER LUIGI ZINZANI

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other
Takeda					х	x	
Roche					x	x	
Eusapharma			x		х	x	
Kyowa Kirin					x	x	
Novartis			x		x	x	
ADC Therap.						x	
Incyte					x	x	
Beigene					x	X	

MOR208: An Enhanced CD19 Antibody

- MOR208 is an Fc-enhanced monoclonal antibody that targets CD19
- Fc-enhancement of MOR208 leads to a potentiation of ADCC and ADCP
- MOR208 induces direct cytotoxicity



Direct cytotoxicity



Katz B-Z et al Leukemia & Lymphoma 2014 Fujimoto M, et al. Immunity 2000

Phase II a: MOR208 in R-R NHL – study design

Multicentre study with 2-stage design (NCT01685008)



Phase II a: MOR208 in R-R NHL – Baseline Characteristics

Characteristic, n (%)		DLBCL n=35	iNHL* n=45	MCL n=12	Total n=92
Age, years	Median	71	66	64.5	66.5
Sex	Male	24 (69)	21 (47)	11 (92)	56 (61)
Ann Arbor stage	1-11	4 (11)	5 (11)	1 (8)	10 (11)
	III-IV	30 (86)	40 (89)	11 (92)	81 (88)
	Missing	1 (3)	0	0	1 (1)
ECOG PS	0-1	34 (97)	43 (96)	11 (92)	88 (96)
	2	1 (3)	2 (4)	1 (8)	4 (4)
Prior lines of therapy	1	12 (34)	16 (36)	3 (25)	31 (34)
	2	8 (23)	6 (13)	1 (8)	15 (16)
	23	15 (43)	23 (51)	8 (67)	46 (50)
Rituximab refractory	Yes	24 (69)	22 (49)	6 (50)	52 (57)
Last rituximab dose	<6 months	14 (40)	6 (13)	1 (8)	21 (23)
Prior stem cell transplantation	Yes	4 (11)	8 (18)	1 (8)	13 (14)

*Includes follicular lymphoma and other indolent NHLs

Data are n (%) unless otherwise stated. Rituximab refractory was defined as patients who demonstrated less than a partial response or response lasting less than 6 months to a prior rituximab-containing regimen

Phase II a: MOR208 in R-R NHL – Best Overall Response Rate



Phase II a: MOR208 in R-R NHL – Duration of Response



Jurczak W and Zinzani PL, Ann Oncol 2018

Phase II a: MOR208 in R-R NHL – PFS



Phase II a: MOR208 in R-R NHL – AE Profile

Grade ≥3 TEAEs,* n (%)	DLBCL n=35	iNHL [†] n=45	MCL n=12	Total n=92
Any [‡]	19 (54)	14 (31)	4 (33)	37 (40)
Hematological [¥]				
Neutropenia	6 (17)	2 (4)	0	8 (9)
Thrombocytopenia	2 (6)	1 (2)	1 (8)	4 (4)
Anemia	3 (9)	0	0	3 (3)
Non-Hematological [¥]				
Dyspnea	2 (6)	1 (2)	1 (8)	4 (4)
Pneumonia	3 (9)	0	0	3 (3)
Fatigue	1 (3)	1 (2)	0	2 (2)
Hypokalemia	1 (3)	1 (2)	0	2 (2)
Infections and Infestations#	5 (14)	1 (2)	0	6 (7)
Infusion-related, n (%)	DLBCL n=35	iNHL [†] n=45	MCL n=12	Total n=92
Any	4 (11)	5 (11)	2 (17)	11 (12)
Grade 1/2	4 (11)	4 (9)	2 (17)	10 (11)
Grade 4	0	1 (2)	0	1 (1)
		There v	were no treatmen	t-related deaths

MOR208 Single Agent in R/R NHL



L-MIND: Trial design



- *Primary refractory is defined as no response to, or progression/relapse during or within 6 months of frontline therapy.
- [†]A loading dose of tafasitamab was administered on Day 4 of Cycle 1. ASCT, autologous stem cell transplantation; DLBCL, diffuse large B-cell lymphoma; DoR, duration of response; ECOG PS, Eastern Cooperative Oncology Group performance status; HDC, high-dose chemotherapy; IRC,independent review committee; LEN, lenalidomide; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; R/R, relapsed or refractory; SD, stable disease.
 1. Salles G, et al. Lancet Oncol 2020;21(7):978-88.

L-MIND: Patient characteristics

Characteristic	Specification	N=81
Age, years [*]	Median (range)	72 (41-86)
Sov. n (%)	Male	44 (54)
Sex, II (70)	Female	37 (46)
And Arbon stores in $(0)^*$	I-II	20 (25)
Ann Arbor stage, n (%)	III-IV	61 (75)
	0-2	40 (49)
KISK (IPI), N (%)	3-5	41 (51)
	Yes	45 (56)
Elevated LDH, n (%)	No	36 (44)
	Median	2
	1	40 (49)
Prior lines, n (%)*	2	35 (43)
	3	5 (6)
	4	1 (1)

Characteristic	Specification	N=81
Dui	Yes	15 (19) ⁺
Primary refractory, n (%)	No	66 (81)
Define the matrices the new line $\pi / 0/1^*$	Yes	36 (44)
Refractory to previous therapy line, n (%)	No	45 (56)
Dring ASCT $n(\ell)$	Yes	9 (11)
	No	72 (89)
	GCB	37 (46)
Centrolly accessed Hans algorithm)	Non-GCB	20 (25)
(Centrany assessed—nans algorithm)	Unknown	24 (30)

•

*At study entry. [†]Primary refractory patients had a DoR to 1 prior line of therapy of 3–6 months.

• ASCT, autologous stem-cell transplant; GCB, germinal center B-cell-like; IHC, immunohistochemistry; IPI, International Prognostic Index; LDH, lactate dehydrogenase; SCT, stem cell transplant.

1. Salles G, et al. Lancet Oncol 2020;21(7):978-88.

L-MIND: Patient disposition after ≥35 months f-up



Adapted from Maddocks K, et al. Poster presented at ASH 2020. Abstract 3021.

Primary endpoint: Best objective response rate (IRC)

Tafasitamab plus LEN	1 prior treatment (N=40)	≥2 prior treatments (N=40)	Overall (N=80)
Best objective response, n (%)			
CR	19 (47.5)	13 (32.5)	32 (40.0)
PR	8 (20.0)	6 (15.0)	14 (17.5)
SD	7 (17.5)	6 (15)	13 (16.3)
PD	5 (12.5)	8 (20.0)	13 (16.3)
NE*	1 (2.5)	7 (17.5)	8 (10.0)
ORR, n (%) [95% CI] ⁺	27 (67.5) [50.9–81.4]	19 (47.5) [31.5–63.9]	46 (57.5) [45.9–68.5]

 *No valid post-baseline response assessments. [†]Two-sided 95% Clopper-Pearson exact method based on a binomial distribution.CI, confidence interval; CR, complete response; IRC, independent review committee; LEN, lenalidomide; NE, not evaluable; ORR, objective response rate; PD, disease progression; PR, partial response; SD, stable disease.

Nowakowski G, et al. ASCO 2020 (Abstract 8020)

.

DoR by number of prior treatment lines



CI, confidence interval; DoR, duration of response; NR, not reached;

Duell J, ICML 2021

PFS by number of prior treatment lines



	1 prior line (n=40)	≥2 prior lines (n=40)	All patients (n=80)
12-month PFS, % (95% CI)	57.8 (39.9, 72.2)	421 (25.7, 57.7)	50.0 (37.8, 61.0)
18-month PFS, % (95% CI)	51.2 (33.4, 66.5)	42.1 (25.7, 57.7)	46.6 (34.5, 57.9)
24-month PFS, % (95% CI)	47.8 (30.2, 63.4)	38.6 (22.6, 54.5)	43.1 (31.1, 54.5)
36-month PFS, % (95% CI)	47.8 (30.2–63.4)	34.3 (18.6–50.7)	41.1 (29.1, 52.7)

Cl, confidence interval; HR, hazard ratio; NR, not reached; PFS, progression free survival.

Duell J, ICML 2021

OS by number of prior treatment lines



24-month OS, % (95% CI)

36-month OS, % (95% CI)

Cl, confidence interval; CR, complete response; NR, not reached; OS, overall survival; PR, partial response...

Duell J, ICML 2021

57.2 (45.1, 67.5)

47.3 (35.5, 58.2)

46.3 (29.8, 61.3)

34.7 (19.9, 50.1)

67.9 (50.4, 80.3)

59.7 (42.3, 73.5)

Patients alive after 3 years of follow-up



OS, overall survival

Duell J, ICML 2021

Favorable safety profile of tafasitamab + LEN followed by extended tafasitamab monotherapy

• No unexpected toxicities or new safety signals were reported from the updated analyses

Event	All Grades (≥10%)	Grade ≥3 (>1 patient)
	All patients	All patients
	N=81	N=81
Hematologic TEAEs, n (%)		
Neutropenia	41 (50.6)	40 (49.4)
Anemia	30 (37.0)	6 (7.4)
Thrombocytopenia	25 (30.9)	14 (17.3)
Non-hematologic TEAEs, n (%)		
Diarrhea	29 (35.8)	1 (1.2%)
Asthenia	20 (24.7)	2 (2.5%)
Cough	22 (27.2)	1 (1.2%)

TEAE, treatment emergent adverse events.

Duell J, ICML 2021

Summary of hematological and non-hematological TEAEs (any Grade) count by patient-years of exposure to tafasitamab



Extended Tafasitamab monotherapy

LEN, lenalidomide; TEAE, treatment emergent adverse event.

1. Salles G, et al. Lancet Oncol 2020;21(7):978-88.

Summary

- Updated analyses with 3-year follow-up confirm the potential of tafasitamab + LEN followed by tafasitamab monotherapy in achieving durable remission and survival benefit
 - Median DoR was 43.9 months across the full analysis set and was comparable regardless of number of prior therapies
 - Median OS was 33.5 months across the full analysis set and was 47.5 months in patients with one prior therapy
- Tafasitamab plus LEN conferred longer PFS and OS in patients who received with 1 vs ≥2 prior lines of therapy
- Decreased burden of TEAEs during extended tafasitamab monotherapy suggests a good tolerability profile for the extended tafasitamab treatment until progression, leading to durable remission
- These data demonstrate the clinical benefit of this chemotherapy-free regimen as a SoC in patients with R/R DLBCL who are ineligible for ASCT, especially those who have received 1 prior line of treatment

• CR, complete response; LEN, lenalidomide NR, not reached; OS, overall survival; SoC, standard of care; PFS, progression free survival; TEAE, treatment emergent adverse events.