**20 ANNI DI EMATOLOGIA
A TREVISO**

TREVISO | 18-20 NOVEMBRE 2021
Auditorium Fondazione Cassamarca

Le lesioni istopatologiche: dall'infezione al linfoma

Dr. Claudio Agostinelli

UO Emolinfopatologia IRCCS Azienda Ospedaliero-Universitaria di Bologna
Dipartimento di Medicina Specialistica, Diagnostica e Sperimentale, Università di Bologna

Disclosures of Claudio Agostinelli: nessun conflitto di interessi da dichiarare

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other

Disordini linfoproliferativi correlati ad infezioni nel tratto gastroenterico

Linfoma della zona marginale extra-nodale (*Helicobacter pylori*)

Malattia da catene pesanti alpha (*Campilobacter Jejuni*)

Linfoma a cellule T/NK extranodale (EBV)

Linfoma diffuso a grandi cellule B EBV+ NAS

Linfoma plasmoblastico (EBV)

«Primary effusion lymphoma» extracavitario (EBV-HHV8)

Ulcera muco-cutanea EBV+

Disordini linfoproliferativi post trapianto (EBV)

paziente

immunocompetente

possibile

immunodeficienza

paziente

immunocompromesso

MALT in gastrite cronica *HP*

Varbanova Best & Practice Research Clinical Gastroenterology 2014

Gastrite follicolare da HP

CD20

CG

Ki67

chronic inflammation (chronic infection or autoimmune processes) sustained antigenic or auto antigenic stimulation not only triggers polyclonal B-cell proliferation, favours the assembly of aggregates of polyclonal B-lymphocytes but also recruits a series of inflammatory cells, including T-lymphocytes, macrophages and neutrophils, to the site of inflammation.

The inflammatory components may promote the growth of neoplastic B-lymphocytes by either one of two mechanisms: **a direct one**, where, for instance, neutrophils release reactive oxygen species (ROS), which cause a wide range of genetic aberrations

an indirect one, where chronic inflammation sustains and prolongs polyclonal proliferation of B-cells, with the consequent increased risk of double-stranded DNA breaks and translocations owing to the inherent genetic instability of B-cells during somatic hypermutation and class switching recombination

Helicobacter pylori e linfomagenesi

Kuo et al Cancers 2019

The cytotoxin-associated gene A (CagA) deregulates intracellular signaling pathways in B-lymphocytes in tyrosine phosphorylation-dependent and -independent manners to initiate the lymphomagenesis of Helicobacter pylori (HP)-dependent gastric mucosa-associated lymphoid tissue (MALT) lymphoma.

Linfoma della zona marginale extranodale

Giemsa

Ki67

CD20

lambda

kappa

CD23⁺ in FDC

Architettura:
multi-focalità,
crescita:
 perifollicolare
 colonizzazione dei follicoli
 diffusa

Citologia

Lymphoepithelial Lesions (LEL)

Histology score to differentiate gastric MALT lymphoma from chronic gastritis

score	interpretation	histology
0	normal	occasional plasmacells
1	chronic active gastritis	lymphocyte clusters, no follicles
2	follicular gastritis	prominent follicles, no lymphoepithelial lesions
3	suspicious, probably reactive	follicles, occasional adjacent lymphoepithelial lesions, no diffuse infiltrate
4	suspicious, probably lymphoma	follicles, diffuse marginal-zone cell infiltrate no lymphoepithelial lesions
5	MALT lymphoma	follicles, diffuse marginal-zone cell infiltrate lymphoepithelial lesions present

Score 3

1° biopsia

Score 3/4

CD20

Score 5: linfoma MALT

CD20

Score 5: linfoma MALT

Fenotipo di esclusione

Follicolare	Mantellare	HCL	LLC
CD10+	CD5+	Annessina+	CD20 w+
BCL6+	Ciclina D1+	Ciclina D1+	CD5+
BCL2+	SOX11+	CD103+	CD23+
LMO2+		CD123+	LEF1+
HGAL+			IRF4 foc+

IRTA1 is selectively expressed in nodal and extranodal marginal zone lymphomas

Brunangelo Falini, Claudio Agostinelli,¹ Barbara Bigerna, Alessandra Pucciarini, Roberta Pacini, Alessia Tabarrini, Flavio Falcinelli, Milena Piccioli,¹ Marco Paulli,² Marcello Gambacorta,³ Maurilio Ponzoni,⁴ Enrico Tiacchi, Stefano Ascani,⁵ Maria Paola Martelli, Riccardo Dalla Favera,⁶ Harald Stein⁷ & Stefano A Pileri¹

Table 1. Immunoglobulin superfamily receptor translocation-associated 1 (IRTA1) expression in 2104 lymphomas

Lymphoid tumours	n	IRTA1 ⁺ cases	%
Chronic lymphocytic leukaemia/SLL	325	0*	0
Lymphoplasmacytic lymphoma	30	0	0
B-cell marginal zone lymphoma			
Splenic	21	0*	0
Nodal	210	154	73
Extranodal	329	307	93
NOS	30	22	73
Follicular lymphoma			
Grade 1/2	130	0*	0
Grade 3A	89	0*	0
Grade 3B	21	0*	0
NOS ⁺	80	0*	0
Mantle cell lymphoma	121	0*	0
Hairy cell leukaemia	30	0	0
Diffuse large B-cell lymphoma	256	69	27
Burkitt lymphoma	71	0	0
Extramedullary plasmacytoma	6	0	0
Classical Hodgkin lymphoma	91	0	0
NLPHL	6	0	0
Peripheral T-cell lymphoma/NOS	160	0	0
Angioimmunoblastic T-cell lymphoma	48	0	0
Anaplastic large-cell lymphoma ALK ⁺	27	0	0
Anaplastic large-cell lymphoma ALK ⁻	17	0	0
EATL	1	0	0
Mycosis fungoides	5	0	0

Myeloid Cell Nuclear Differentiation Antigen (MNDA)

IRTA1 and MNDA Expression in Marginal Zone Lymphoma

Utility in Differential Diagnosis and Implications for Classification

Zhen Wang, MD, PhD, and James R. Cook, MD, PhD

From the Department of Laboratory Medicine, Robert J. Tomsich Pathology and Laboratory Medicine Institute, Cleveland Clinic, Cleveland, OH.

Key Words: IRTA1; MNDA; Marginal zone lymphoma

Am J Clin Pathol March 2019;151:337-343

Modalità di diffusione

- ENMZL gastrici tendono a localizzarsi in strutture mucose
- diffusione intragastrica e intestino (per interazione integrine $\alpha 4\beta 7$ con HEV)
- 25% dei ENMZL gastrici hanno diffusione subclinica ad altre mucose (50% in ENMZL di altre sedi)
- la disseminazione secondaria allo stomaco è più rara in genere da ENMZL polmonari o intestinali (gastroscopia di staging)

ENMZL polmone

Profondità infiltrato:

mucosa/sottomucosa = regr.⁺

muscolare propria = regr.⁻

eco-endoscopia.

Valutazione regressione:

istologia (più attendibile di PCR);

tempo richiesto: 2-28 mesi.

La regressione può essere rapida (4 mesi) ma in buona parte dei casi può ritardare fino a 1 o > anni

Table I. GELA histological grading system for post-treatment evaluation of gastric MALT lymphoma.

Score	<u>Lymphoid infiltrate</u>	<u>LEL</u>	<u>Stromal changes</u>
Complete Histological remission (CR)	Absent or scattered plasma cells and small lymphoid cells in the LP	Absent	Normal or empty LP and/or fibrosis
Probable minimal residual disease (pMRD)	Aggregates of lymphoid cells or lymphoid nodules in the LP/MM and/or SM	Absent	Empty LP and/or fibrosis
Responding residual disease (rRD)	Dense, diffuse or nodular, extending around glands in the LP	Focal LEL or absent	Focal empty LP and/or fibrosis
No change (NC)	Dense diffuse or nodular	Present 'may be absent'	No changes

In follow-up:

- HP è carcinogeno di classe I (IARC) per il carcinoma gastrico
- pazienti con G-ENMZL sono a maggior rischio di carcinomi gastrici metacroni
- riscontro di metaplasia intestinale in 50-70% dei pazienti a 6-18 mesi dalla diagnosi con G-EMZL
- Per quanto tempo mantenere follow-up? (rischio di recidive tardive di linfoma e rischio sviluppo carcinoma)

Ahn HJ *et al.* *H. pylori* in gastric carcinogenesis

World J Gastroenterology 2015

Paz. M 71aa

ENMZL: alterazioni genomiche numeriche e strutturali

Gains / Losses (20-40%, > che in altri linfomi):

+18/18q ricorrente

+3/3q ricorrente

del 6q23.3 (nel 15-30% TNFAIP3/A20)

Traslocazioni:

t(11;18)(q21;q21) *BIRC3/MALT1*

t(1;14)(p22;q32) *IGH/BCL10*

t(14;18)(q32;q21) *IGH/MALT1*

t(3;14)(p13;q32) *IGH/FOXP1*

(Isaacson, Nat Rev Cancer 2004; 4:644-53)

Nature Reviews | Cancer

Frequenza diversa in base alla sede

Site	t(11;18) (q21;q21)	t(14;18) (q32;q21)	t(3;14) (p14.1;q32)	t(1;14) (p22;q32)	+3	+18
Stomach	6-26	1-5	0	0	11	6
Intestine	12-56				75	25
Ocular adnexa/orbit	0-10				38	13
Salivary glands	0-5				55	19
Lung	31-53				20	7
Skin	0-8	0-14	0-10	0	20	4
Thyroid	0-17	0	0-50	0	17	0

t(11;18)(q21;q21):

- Minore risposta alla terapia eradicante HP
- Malattia più estesa
- Progression free survival più corta 26% vs 57%

¹Data summarized according to Streubel *et al.* {2109A} and Remstein *et al.* {1836}.

Certain translocations are associated with particular sites of origin (WHO 2008)

TBL1XR1 24%
 GPR34 16%
 NOTCH2 11%
 SPEN 11%
 KMT2C 11%

TET2 65%
 TNFRSF14 44%
 PIK3CD 23%
 CCR6 8%
 FOXO1 8%

KMT2D 25%
 TNFAIP3 18%
 PRDM1 12%
 NOTCH1 12%
 EP300 11%

KLF2 18%
 NOTCH2 16%
 TP53 12%
 TNFAIP3 8%
 KMT2D 7%

FAS 63%
 SLAMF1 24%
 SPEN 18%
 NCOR2 13%
 CASP10 11%

Salivary gland MZL

Thyroid MZL

Pulmonary MZL

Splenic MZL

Cutaneous MZL

Dural MZL

Ocular adnexal MZL

Gastric MZL

Nodal MZL

TNFAIP3 45%
 NOTCH2 36%
 TLBXR1 36%
 EP300 18%
 KLHL6 18%

TNFAIP3 32%
 KMT2D 15%
 KMT2C 11%
 LRP1B 11%
 MYD88 10%

NOTCH1 16%
 TNFAIP3 15%
 NF1 14%
 TRAF3 13%
 TET2 11%

KMT2D 20%
 TNFAIP3 14%
 CREBBP 12%
 PTPRD 12%
 KLF2 12%

Vela et al Virchows Archive 2021

EMZL gastrico H. pylori negativo

- Aumento dei casi di ENMZL gastrici HP- in ultimi 20 aa:
da 5%-10% a 30-50%
- La persistenza della flogosi cronica può portare a scomparsa istologia di HP ma breath-test, sierologia e test feci+
- Spiegazioni:
test falsi negativi
infezioni da altri batteri (*H. heilmannii*, *H. suis*, *H. bizzozeri*, *salomnois*)?
associazione con malattie autoimmuni (tiroiditi di Hashimoto)?

Tessuto MALT in tiroidite di Hashimoto

Genetic Characterization and Clinical Features of *Helicobacter pylori* Negative Gastric Mucosa-Associated Lymphoid Tissue Lymphoma

Barbara Kiesewetter ^{1,2,*}, Christiane Copie-Bergman ³, Michael Levy ⁴, Fangtian Wu ^{1,5}, Jehan Dupuis ⁶, Caroline Barau ⁷, Luca Arcaini ^{8,9}, Marco Paulli ¹⁰, Marco Lucioni ¹⁰, Arturo Bonometti ¹¹, Antonio Salar ¹², Concepción Fernández-Rodríguez ¹³, Miguel A. Piris ¹⁴, Francesco Cucco ¹, Rachel Dobson ¹, Yan Li ^{1,15}, Zi Chen ¹, Cyrielle Robe ³, Ingrid Simonitsch-Klupp ¹⁶, Andrew Wotherspoon ¹⁷, Markus Raderer ² and Ming Qing Du ^{1,*}

Cancers 2021

In G-ENMZL HP- :

t(11;18)(q21;q21)

e iper-espressione BCL10

- riarrangiamenti di MALT1 (40%)
- Riarrangiamenti IGH (verosimili t1;14)
- Mutazioni di attivanti NF-kB mutualmente esclusive con traslocazioni *MALT1*
- In più dell' 85% dei casi attivazione *NF-kB*

Linfoma marginale
del piccolo intestino
(Paz M 40aa con
perforazione)

ALPHA HEAVY CHIAN DISEASE

- ❖ (IPSID: Immuno-Proliferative Small Intestine Disease; linfoma medi-terraneo)
- ❖ LZM a marcatissima differenziazione plasmacellulare (DD plasmocitoma extramidollare)
- ❖ Raro; area mediterraneo, a basso livello socio-economico, con infezioni GI frequenti
- ❖ Pazienti II-III decade
- ❖ Intestino: duodeno > digiuno > ileo
- ❖ sindrome da malassorbimento, diarrea intermittente e dolore addominale della durata da alcuni mesi ad anni, episodi occlusione intestinale

Mutazioni puntiformi delezioni/inserzioni durante processo di ipermutazione somatica con conseguente difettiva sintesi della CH1 delle catene pesanti α

α HC in assenza di catena leggera in Siero, urine e fluidi corporei

Elettroforesi sierica normale, riscontro α HC patologica con immunofissazione e immunoelettroforesi

Bianchi Current Gastroenterology report 2018

Campylobacter jejuni:

- ❖ Gram-negativo a forma di elica
- ❖ rappresenta una delle cause più comuni di gastroenterite nel mondo
- ❖ ipotesi patogenetica: Infezioni ricorrenti inducono stato di immunoparesi che impedisce clearance CJ e lo stimolo Ag cronico porta accumulo pc IgA
- ❖ danno genomico può indurre α HCD
- ❖ assenza traslocazioni del EMZL
- ❖ Predisposizione individuale (specifici HLA o gruppi sanguigni)

Disordini linfoproliferativi correlati ad infezioni nel tratto gastroenterico

Linfoma della zona marginale extra-nodale (*Helicobacter pylori*)

Malattia da catene pesanti alpha (*Campilobacter Jejuni*)

Linfoma a cellule T/NK extranodale (EBV)

Linfoma diffuso a grandi cellule B EBV+ NAS

Linfoma plasmoblastico (EBV)

«Primary effusion lymphoma» extracavitario (EBV-HHV8)

Ulcera muco-cutanea EBV+

Disordini linfoproliferativi post trapianto (EBV)

paziente

immunocompetente

possibile

immunodeficienza

paziente

immunocompromesso

Germinal centre model

EBV

a EBV genome: latent genes

Table 2. EBV-related proteins expressed in various diseases according to the latency patterns

Latency Pattern	EBV-Associated Proteins			Disorder
	EBNA1/LMP2A	LMP1/LMP2B	EBNA2/3A/3B/3C/LP	
I	+	-	-	Burkitt lymphoma
II	+	+	-	Classic Hodgkin lymphoma, PEL, AITL, NK/T-cell lymphoma, nasopharyngeal carcinoma, EBV-positive DLBCL of the elderly
III	+	+	+	PTLD, AIDS-related lymphoma, infectious mononucleosis, EBV-positive DLBCL of the elderly

Abbreviations: AIDS, acquired immunodeficiency syndrome; AITL, angioimmunoblastic lymphoma; DLBCL, diffuse large B-cell lymphoma; EBNA, EBV nuclear antigen; EBV, Epstein-Barr virus; LMP, latent membrane protein; LP, leader protein; PEL, primary effusion lymphoma; PTLD, post-transplant lymphoproliferative disorder.

EBV positive Mucocutaneous Ulcer (EBVMCU)

Published in final edited form as:

Am J Surg Pathol. 2010 March ; 34(3): 405–417. doi:10.1097/PAS.0b013e3181cf8622.

EBV Positive Mucocutaneous Ulcer – A study of 26 cases associated with various sources of immunosuppression

Stefan D Dojcinov, M.D., F.R.C.Path⁽¹⁾, Girish Venkataraman, M.D.⁽²⁾, Mark Raffeld, M.D.⁽²⁾, Stefania Pittaluga, M.D., Ph.D.⁽²⁾, and Elaine S Jaffe, M.D.⁽²⁾

Entità provvisoria
nella classificazione WHO 2017;
non è considerato linfoma
ma disordine linfoproliferativo

Mature B-cell neoplasms
Chronic lymphocytic leukemia/small lymphocytic lymphoma
Monoclonal B-cell lymphocytosis*
B-cell prolymphocytic leukemia
Splenic marginal zone lymphoma
Hairy cell leukemia
<i>Splenic B-cell lymphoma/leukemia, unclassifiable</i>
<i>Splenic diffuse red pulp small B-cell lymphoma</i>
<i>Hairy cell leukemia-variant</i>
Lymphoplasmacytic lymphoma
Waldenström macroglobulinemia
Monoclonal gammopathy of undetermined significance (MGUS), IgM*
μ heavy-chain disease
γ heavy-chain disease
α heavy-chain disease
Monoclonal gammopathy of undetermined significance (MGUS), IgG/A*
Plasma cell myeloma
Solitary plasmacytoma of bone
Extrasosseous plasmacytoma
Monoclonal immunoglobulin deposition diseases*
Extranodal marginal zone lymphoma of mucosa-associated lymphoid tissue (MALT lymphoma)
Nodal marginal zone lymphoma
<i>Pediatric nodal marginal zone lymphoma</i>
Follicular lymphoma
In situ follicular neoplasia*
Duodenal-type follicular lymphoma*
Pediatric-type follicular lymphoma*
<i>Large B-cell lymphoma with IRF4 rearrangement*</i>
Primary cutaneous follicle center lymphoma
Mantle cell lymphoma
In situ mantle cell neoplasia*
Diffuse large B-cell lymphoma (DLBCL), NOS
Germinal center B-cell type*
Activated B-cell type*
T-cell/histiocyte-rich large B-cell lymphoma
Primary DLBCL of the central nervous system (CNS)
Primary cutaneous DLBCL, leg type
EBV ⁻ DLBCL, NOS*
<i>EBV⁺ mucocutaneous ulcer*</i>
DLBCL associated with chronic inflammation
Lymphomatoid granulomatosis
Primary mediastinal (thymic) large B-cell lymphoma
Intravascular large B-cell lymphoma
ALK ⁺ large B-cell lymphoma
Plasmablastic lymphoma
Primary effusion lymphoma
<i>HHV8⁺ DLBCL, NOS*</i>

EBVUMC

- Ulcere solitarie, a margini netti, a lenta crescita
- sede: orofaringea (50%), cute (30%), GI (20%)
- mai formazione di masse, linfoadenopatie generalizzate, organomegalie o coinvolgimento osteomidollare
- correlate a immunosoppressione iatrogena:
metotrexate, ciclosporina, azatioprina, prednisolone
tacrolimus, inibitori TNF α (infliximab), micofenolato mofetile
- correlazione con immunosenescenza (pazienti anziani), HIV

Nel tratto GI in pazienti:

- immunosoppressi per trattamento di IBD (Crohn, RCU)
- artrite reumatoide
- post-trapianto
- trattati per linfoma
- coliti immunorelate (post trattamento ICPI anti CTLA4 /PD1)
- diverticolite

EBVMCU è caratterizzata da un decorso autolimitante con remissioni spontanee o regressione dopo riduzione o sospensione della terapia immunosoppressiva

(Ishigava Cancers 2021, Zanelli Cancers 2021)

Ulcera ileale

- Grandi cellule sparse citologia immunoblastica o simil Reed-sternberg, o in apoptosi
- piccoli linfociti, eosinofili, istiociti, plasmacellule
- rima di piccoli linfociti alla base

CD3

CD30

- I basti esprimono in grado variabile CD20
- CD30⁺ ± CD15⁺
- PAX5⁺, IRF4⁺

Fenotipo simil Hodgkiniano

- piccoli linfociti CD4⁺ e CD8⁺ (spesso attivati)

- EBV+ in piccole grandi cellule
- 30% *IG* monoclonali
- Possibile riscontro mono/oligo cloni T:
riduzione della diversità del repertorio del
TCR legata alla senescenza

EBVMCU: prognosi e terapia

- Andamento indolente autolimitante
- in caso di immunosoppressione iatrogena
una riduzione/interruzione del farmaco induce remissione
- possibile andamento con remissioni/recidive
- EBVMCU in post-linfoma e IBD possibili ulcere multiple
- in coliti immunorelate possibili perforazioni
- raramente evoluzione a linfoma (in IBD)
- possibile terapia: chirurgica, anti CD20/anti CD30,
radioterapia (o combinazioni)

Attenzione alla diagnosi differenziale

lesione del sigma CD30+

EBVMCU vs EBV+DLBCL

EBVMCU vs Hodgkin classico

- Paziente di 39 aa in terapia per RCU
- EBVMCU non forma masse

Disordine linfoproliferativo
associato a immunospressione
iatrogena tipo cHL