

gli under 40 a confronto

Verona, Centro Congressi Camera di Commercio 26-27 settembre 2025

Ruolo della re-biopsy: aspetti clinici

Tetiana Skrypets

Ematologia, Istituto Tumori "Giovanni Paolo II", Bari, Italy


Disclosures of Name Surname

Company name	Research support	Employee	Consultant	Stockholder	Speakers bureau	Advisory board	Other
Abbvie					X		X
BeONE					X		X
J&J					X		X
Eli Lilly							X
Menarini					X		
Novartis	X						
Takeda					X	X	X


AGENDA


- 1. Importance of accurate diagnosis and disease monitoring
- 2. Rationale and clinical indications for rebiopsy
- 3. Practical Considerations
- 4. Emerging Technologies and Future Directions
- 5. Summary and Conclusions


Importance of accurate diagnosis and disease monitoring




- 1. Kaplan-Meier survival estimate for patients with concordant versus discordant pathologic diagnosis after MGH review with number of patients at risk and 95% confidence limits. Massachusetts General Hospital. Chipidza et al., JCO Glob Oncol, 2021
- 2. Kaplan-Meier plots of EFS and OS of MGZL and PMBL. **Wyndham et al., Blood, 2014**
- 3. Kaplan-Meier plots of OS of PTCLs and HL. ICML 2025, poster 388 Luminari et al. Ansel et al., N Engl J Med, 2022











Rationale and clinical indications for re-biopsy


- 1. Heterogeneity and evolution of lymphoma at molecular and histologic levels
- 2. Limitations of initial biopsy and diagnosis
- 3. Changes in disease biology over time due to therapy or progression
- 4. Need for updated tissue diagnosis to guide treatment decisions
- 5. Suspected disease relapse or progression
- 6. Transformation to a more aggressive lymphoma subtype (e.g., FL to DLBCL, Richter)
- 7. Discrepant clinical or imaging findings
- 8. Assessment before initiating novel therapies (e.g., CAR-T, targeted agents)
- 9. Evaluation of resistance mechanisms post-therapy


Heterogeneity and evolution of lymphoma at molecular and histologic levels






- 1. R. Kansal et al, *Lymphatics* 2023, 1(2), 155-219
- Sánchez-Beato et al, Clinical and Translational Oncology (2024) 26:1043–1062
- 3. Sánchez-Beato et al, Clinical and Translational Oncology (2024) 26:1043–1062
- Daniel J. Landsburg, MD, and Brad S. Kahl, MD, https://dailynews.ascopubs.org/do/exploiting-molecular-heterogeneity-aggressive-b-cell-lymphomas-develop-effective-new







They are histologically classified as Hodgkin or non-Hodgkin lymphoma (NHL), with NHL accounting for 90% of cases

Summary of genetic alterations in B/T-cell lymphomas and their clinical utility

B-cell Lymphoma		Genetic alterations	Clinical significance	
Burkitt Lymphoma (BL)	EBV+	MYC rear., MYC aSHM, CDKN2A, DDX3X	DIAGNOSTIC	
		TP53	PROGNOSTIC	
	EBV -	MYC rear., TCF3, ID3, CDKN2A, DDX3X	DIAGNOSTIC	
		TP53	PROGNOSTIC	
Hodgkin Lymphoma (HL)		XPO1, EP300, CREBBP, TP53, B2M, NFKBIE, TNFAIP3, STAT3, STAT6, PTPN1, ITPKB, GNA13, ARID1A, KTM2D, IGLL5, CSFR2B, BTK	FUTURE	
Diffuse large B-cell lymphoma, not otherwise s NOS)	pecified (DLBCL-	NOTCH2, BCL10, TNFAIP3, UBE2A, CD70, CCND3, DTX1, BCL2, EZH2, CREBBP, TNFRSF14, KMT2D, IRF8, EP300, GNA13, MYD88, CD79B, PIM1, PIM2, PRDM1, BTG1, CD58, NOTCH1, SGK1, SOCS1, TET2, STAT3, TP53; MYC rear., BCL2 rearr., BCL6 rear	DIAGNOSTIC/PROGNOSTIC	
Large B-cell lymphoma with IRF4 rearrangeme	ent	IRF4 rear, IRF4, CARD11, MYD88, CD79B	DIAGNOSTIC	
Large B-cell lymphoma with 11q aberration		11q aberration, GNA13	DIAGNOSTIC	
High grade B-cell lymphoma (HGBCL)	HGBCL-DH-BCL2	MYC rear., BCL2 rear., BCL2, KMT2D, CREBBP, TNFRS14, EZH2	DIAGNOSTIC/PROGNOSTIC	
	HGBCL-DH-BCL6	MYC rear., BCL6 rear	DIAGNOSTIC	
	HGBCL-NOS	MYD88, CD79B, TBL1XR1, TP53, KMT2D	DIAGNOSTIC	

Sánchez-Beato et al, Clin and Transl Oncology (2024)

Summary of genetic alterations in B/T-cell lymphomas and their clinical utility

B-cell Lymphoma		Genetic alterations	Clinical significance
Chronic lymphocytic leukemia (CLL)		IGVH SHM, Del(17p)/TP53, Del(11q), Del(13q), trisomy 12, NOTCH1, SF3B1, ATM, BIRC3, NFKBIE, EGR2, MYD88, XPO1, CHD2	DIAGNOSTIC/PROGNOSTIC
		BTK, PLCG2, BCL2, TP53, CARD11	PREDICTIVE
Lymphoplasmacytic lymphoma (LPL)		MYD88 ^{L265P} #, CXCR4	DIAGNOSTIC/PROGNOSTIC/PREDICTIVE
Marginal zone lymphoma (MZL)	Spleenic MZL	KLF2, NOTCH2, TP53, NOTCH1, MLL2, ARID1A, SIN3A, TNFAIP3, MYD88, CARD11, trisomies 3, 18, Del(7q)	DIAGNOSTIC
	Nodal MZL	<i>KLF2, NOTCH2, KMT2D, PTPRP,</i> trisomies <i>3, 18, 7, 12</i>	DIAGNOSTIC
	Extranodal MZL	trisomies 3, 18, 12, T(11;18)#, T(1;14), T(3;14), T(14;18) (IGH::MALT1); TNFAIP3, CD79A, CD79B, CARD11, BIRC3, TRAF3, TNFRSF11A	DIAGNOSTIC/PROGNOSTIC (m7-FLIPI)/ THERAPY
Follicular lymphoma (FL)	FL	T(14;18); KMT2D, EZH2*#, CREBBP#, EP300#, MEF2B#, ARID1A#, FOXO1#, CARD11#	DIAGNOSIS/PROGNOSTIC#/THERAPY*
		NOTCH2, DTX1, UBE2A, HIST1H1E, MYC, TP53, CCND3, GNA13, S1PR2, P2RY8, POU2AF1, CDKN2A/B loss	PROGNOSTIC (HT)
	Diffuse FL	TNFRSF14, STAT6, CREBBP, EZH2	DIAGNOSTIC
	Pediatric TFL	TNFRSF14, MAP2K1	DIAGNOSTIC
	Duodenal type FL	TNFRSF14, CREBBP, EZH2	DIAGNOSTIC
Mantle cell lymphoma (MCL)		T(11;14), CCND2 and CCND3 rearr	DIAGNOSTIC
		IGVH SHM, Del(17p)/TP53, ATM, NOTCH1/2, KMT2D	PROGNOSTIC
		Del(17p)/TP53, BIRC3, TRAF2, NSD2, CARD11	PREDICTIVE Sánchez-Beato

T-cell Lymphoma		Genetic alterations	Clinical significance
Follicular helper T-cell lym-phoma (TFH)	Angioimmunoblastic (AITL) Follicular type Not otherwise specified (NOS)	IDH2 ^{R172} , RHOA ^{G17V} , TET2, DNMT3A\$, VAV1, CD28, ICOS, FYN and LCK rearr	DIAGNOSTIC/PREDIC- TIVE ^{\$}
Anaplastic large cell lym-	ALK-positive	ALK fusion, NOTCH1*	DIAGNOSTIC/THERAPY*
phoma (ALCL)	ALK-negative	DUSP22 rearr#, JAK1, JAK3, STAT3 Del(17p)/TP53, TP63 rearr., PRDM1 loss	DIAGNOSTIC/PROGNOS- TIC*/PREDICTIVE PROGNOSTIC
Peripheral T-cell lymphoma, not otherwise specified (PTCL-NOS)	PTCL-TBX21 PTCL-GATA3	TET1, TET3, DNMT3A TP53#, PRDM1, CDKN2A/B, RB1 and PTEN loss, STAT3 and MYC gain	DIAGNOSTIC/PROGNOS- TIC#
Primary nodal EBV-positive T	T-/NK-cell lymphoma	TET2, PI3KCD, STAT3, TP53, CARD11	DIAGNOSTIC
Cutaneous T-cell lymphoma (CTCL)		PLCG1, NFATC2, NFAT5, ZEB1, PRKCQ, RHOA, VAV1, PREX2, CTCF, ARID1A, TRRAP	DIAGNOSTIC
Hepatosplenic T-cell lymphoma (HTCL)		STAT5B, STAT3, PIK3CD, SETD2, IN080, ARID1; Loss of 7p, amplification of 7q	DIAGNOSTIC
Breast implant-associated ALCL		STAT3, JAK1, JAK3, DNMT3, TP53	DIAGNOSTIC
Extranodal NK/T-cell lymphoma (ENKT), nasal type		TP53#, DDX3X#, Del(6q), STAT3, JAK3, STAT5B	PROGNOSTIC [#] Sánchez-Bea

Why Re-Biopsy? The Clinical Rationale

Key Clinical Questions to Address	Answer, example			
Is it a true relapse?	Imaging (PET/CT scans): - show metabolic activity, <u>BUT NOT</u> always definitively differentiate between lymphoma, inflammation, false-positive or other conditions (e.g., sarcoidosis) -biopsy provides a <u>definitive histological confirmation</u> .			
- particularly important for indolent lymphomas (FL) - transform into a more aggressive subtype (DLBCL) "histologic transformation" requires a completely difference more intensive treatment approach				
Has the molecular and genetic profile evolved?	 lymphoma's <u>biology may change</u> over time <u>new mutations or genetic alterations</u> may emerge <u>affect</u> prognosis and treatment options. 			

Clinical Scenarios for Re-Biopsy

Scenario 1: Suspected Relapse/Progression

- ➤ A patient on follow-up or after completing therapy shows new or growing lesions on a physical exam or imaging (PET/CT, CT scan)
- Crucial when imaging findings are equivocal or a different diagnosis is in the differential.

Scenario 2: Histologic Transformation

- ➤ Patient with indolent lymphoma (FL, MZL) has a sudden change in clinical course:
- ✓ Rapidly enlarging I/nodes or masses
- ✓ Development of B-symptoms
- ✓ Elevated serum LDH...

A re-biopsy is mandatory

Scenario 3: Refractory Disease

- ➤ A patient is not responding to therapy
- Re-biopsy can help to identify resistance mechanisms or lymphoma's biology that may guide the selection of a new, more effective therapy...?
- Another diagnosis (different lymphoma subtype...)

QUESTIONABLE


Articles

Repeat biopsy in relapsed or refractory diffuse large B cell lymphoma: a nationwide survey and retrospective study

Tamar Berger 🚾 📵, Karyn Revital Geiger, Moshe Yeshurun, Anat Gafter-Gvili, Tzippy Shochat, Ronit Gurion,


Pia Raanani & Oren Pasvolsky D ...show less
Pages 2461-2468 | Received 02 Apr 2022, Accepted 18 May 2022, Published online: 28 May 2022

Leukemia & Lymphoma



Reasons for avoiding a biopsy in suspected R/R disease


- > A two-part study: a nationwide case-vignette survey
- ▶ 64 participating physicians opted not to re-biopsy in at least one scenario, more often in refractory cases.
- ➤ In the **retrospective part**, 116 R/R aNHL cases among 61 pts were identified
- Re-biopsy not performed in 72% more in refractory mostly due to low likelihood of alternative diagnoses or problematic location for biopsy.



Reasons for avoiding a biopsy in R/R DLBCL – stratified by line of therapy


PET/CT-guided re-biopsy may find clinically unsuspicious transformation of FL

- > Retrospective study
- ➤ Pts who had undergone PET/CT during primary diagnosis or relapse of FL between 2010 and 2020
- The diagnosis changed from an indolent disease to a transformed lymphoma in >10% (7/63)
- ➤ The HT risk associated with high SUVmax (>10) was 24% (7 of 29 performed biopsies)
- ➤ 4 out of these 7 pts with verified HT had NO previous clinical suspicion of transformation



A comparison of individual SUVmax values between pts with HT vs no HT, with the sensitivity and specificity for HT of the threshold 26.5.

The graph includes patients with SUVmax <10 and a new biopsy



Therapeutic Implications of Re-Biopsy

Re-biopsy findings have a direct impact on patient management by guiding subsequent therapeutic decisions


Personalized Treatment Planning:

- Histologic Transformation:
- > indolent to aggressive lymphoma
- > change to an intensive regimen
- Molecular Profiling:
- > new mutations or markers

Prognostic Value:


- patient risk-stratification
- disease course
- possible treatment options

SCHOLAR-1 STUDY IN REFRACTORY DLBCL



Crump et al., BLOOD, 2017

Reduced CD20 expression yields inferior survival in patients with B-cell lymphoma treated with CD20×CD3 antibodies




- > 3 Danish centers
- Pts received
 CD20×CD3 bispecifics
- Period 2017 -2024 in phase 1/2 trials
- ➤ 148 pretreatment and 60 post-treatment samples


Survival in strong, reduced, and negative immunohistochemical expression of CD20. (A) PFS of patients with either strong, reduced, or negative IHC expression of CD20 antigen. (B) OS of patients with either strong, reduced, or negative IHC expression of CD20 antigen.

Kyvsgaard, Blood 2025

Challenges and Considerations in Re-Biopsy



- 1. Fitzpatrick et al, Cytopathology, 2024
- 2. Hasserjian, The Hematologist, 2023



Size Matters in Lymphoma Diagnosis Tissue Biopsy

1. Invasiveness and Risk:

Tissue biopsy is an invasive procedure with potential risks, including bleeding and infection.

2. Adequacy of Tissue:

A core needle biopsy may not always provide enough tissue for all necessary tests.


3. Logistical Challenges:

Re-biopsy can delay the initiation of salvage therapy.

Emerging Technologies: The Rise of Liquid Biopsy


Advantages of Liquid Biopsy:


- Minimally invasive and can be easily repeated for longitudinal monitoring.
- Can provide a more comprehensive view of tumor heterogeneity.



Current Role and Future Potential:

- > Used to monitor treatment response and detect minimal residual disease (MRD).
- Complementary tool; not a replacement for tissue biopsy





OR PET EOT repositivized during follow-up

Conclusions: A Balanced Approach

1. Tissue re-biopsy remains a cornerstone of care in specific situations for patients with relapsed or refractory lymphoma.

- > Histological confirmation of relapse.
- > Detection of histologic transformation.
- Guidance for therapeutic decisions based on updated molecular and genetic profiles
- 2. A collaborative, multidisciplinary approach is essential.
- 3. The future of lymphoma management will likely involve a synergistic approach
 - > tissue biopsies provide definitive diagnosis
 - liquid biopsies: dynamic, real-time tool for monitoring and guiding treatment